

2026-03-18
2025-2012

Group	11	Fungicide
-------	----	-----------

DISARM™ 480 SC FUNGICIDE

FOR CONTROL OF DISEASES IN TURF

SUSPENSION

COMMERCIAL

ACTIVE INGREDIENT: fluoxastrobin... 480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0193%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037% as preservatives

REGISTRATION NUMBER 30811 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

POTENTIAL SKIN SENSITIZER

NET CONTENTS: 250 mL to 1,000L

For product information call: 1-800-438-6071

UPL AgroSolutions Canada Inc.
PO Box 12219
Research Triangle Park, NC 27709

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

2026-03-18

2025-2012

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS**KEEP OUT OF REACH OF CHILDREN**

Potential skin sensitizer. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Chemical-resistant gloves should be made of any waterproof material, such as nitrile, butyl, neoprene and/or barrier laminate. These are only some of the glove materials that are chemically resistant to this product.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining equipment. If there are no such instructions for washables, use detergent and hot water. Keep and wash clothing separately from other laundry.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Do not enter or allow entry into treated turf until residues are dry.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

The residues of fluoxastrobin are persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing fluoxastrobin during the previous year.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland. Care should be taken when using this product in an integrated pest management program where users are relying on the presence of beneficial arthropods.

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, or other sensitive areas that may be exposed to spray drift.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides and fertilizers. Store this product away from food and feed, in its original container and out of reach of children, preferably in a locked storage area. DISARM 480 SC Fungicide is not affected by freezing.

SPILL CLEAN-UP: Follow safety precautions as directed for handling the product. If on a floor or hard surface, wash the surface or floor with detergent and water, then rinse. If on soil, collect surface soil contaminated with the product.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial or territorial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty rinsed container unsuitable for further use.
4. Dispose of the container in accordance with provincial or territorial requirements.
5. For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

2026-03-18
2025-2012

Group	11	Fungicide
-------	-----------	-----------

DISARM™ 480 SC FUNGICIDE

FOR CONTROL OF DISEASES IN TURF

SUSPENSION

COMMERCIAL

ACTIVE INGREDIENT: fluoxastrobin... 480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0193%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037% as preservatives

REGISTRATION NUMBER 30811 PEST CONTROL PRODUCTS ACT

**READ THE LABEL AND ATTACHED BOOKLET BEFORE
USING**

KEEP OUT OF REACH OF CHILDREN

POTENTIAL SKIN SENSITIZER

NET CONTENTS: 250 mL – 1,000 L

For product information call: 1-800-438-6071

UPL AgroSolutions Canada Inc.
PO Box 12219
Research Triangle Park, NC 27709

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

2026-03-18

2025-2012

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS**KEEP OUT OF REACH OF CHILDREN**

Potential skin sensitizer. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Chemical-resistant gloves should be made of any waterproof material, such as nitrile, butyl, neoprene and/or barrier laminate. These are only some of the glove materials that are chemically resistant to this product.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining equipment. If there are no such instructions for washables, use detergent and hot water. Keep and wash clothing separately from other laundry.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Do not enter or allow entry into treated turf until residues are dry.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

The residues of fluoxastrobin are persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing fluoxastrobin during the previous year.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland. Care should be taken when using this product in an integrated pest management program where users are relying on the presence of beneficial arthropods.

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, or other sensitive areas that may be exposed to spray drift.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides and fertilizers. Store this product away from food or feed, in its original container and out of reach of children, preferably in a locked storage area. DISARM 480 SC Fungicide is not affected by freezing.

SPILL CLEAN-UP: Follow safety precautions as directed for handling the product. If on a floor or hard surface, wash the surface or floor with detergent and water, then rinse. If on soil, collect surface soil contaminated with the product.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial or territorial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty rinsed container unsuitable for further use.
4. Dispose of the container in accordance with provincial or territorial requirements.
5. For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

GENERAL INFORMATION

DISARM™ 480 SC Fungicide is a broad-spectrum fungicide for the control of diseases in turf. DISARM 480 SC Fungicide works by interfering with respiration in plant- pathogenic fungi, and is a potent inhibitor of spore germination and mycelial growth.

DIRECTIONS FOR USE

For use to control disease in turf on sod farms, golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Keep children and pets out of treated area until spray has dried.

DO NOT apply by air.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

UNDER CERTAIN CONDITIONS CONDUCIVE TO EXTENDED INFECTION PERIODS, ADDITIONAL FUNGICIDE APPLICATIONS BEYOND THE NUMBER ALLOWED BY THIS LABEL MAY BE NEEDED. UNDER THESE CONDITIONS, USE ANOTHER FUNGICIDE FROM A DIFFERENT MODE OF ACTION GROUP REGISTERED FOR THE DISEASE.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that DISARM 480 SC Fungicide contains a Group 11 fungicide. Any fungal population may contain individuals naturally resistant to DISARM 480 SC Fungicide and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of DISARM 480 SC Fungicide or other Group 11 fungicides with different groups that control the same pathogens.
- For control of pythium blight and anthracnose, do not apply more than 1 application of Disarm 480 SC Fungicide before alternating with a fungicide having a different mode of action registered for the same disease or pathogen. For control of diseases other than pythium blight and anthracnose, do not apply more than 2 sequential applications before alternating with a fungicide having a different mode of action registered for the same disease or pathogen.

- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify UPL AgroSolutions Canada Inc at 1-800-438-6071 if reduced sensitivity of the pathogen to DISARM 480 SC FUNGICIDE is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact UPL AgroSolutions Canada Inc at 1-800-438-6071.

APPLICATION GUIDELINES

Broadcast Ground Sprayers

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use. Use a pump with the capacity to: (1) maintain a minimum of 240 kPa at nozzles, and (2) provide sufficient agitation in the tank to keep the mixture in suspension (this requires recirculation of 10% of the tank volume per minute). Use jet agitators or a liquid sparge tube for vigorous agitation. Use screens to protect the pump and to prevent nozzles from clogging.

Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh screens at the nozzles. Check nozzle manufacturer's recommendations. For information on spray equipment and calibration, consult sprayer manufacturer's and/or provincial or territorial recommendations. For specific local directions and spray schedules, consult the current provincial or territorial recommendations.

Ground Boom Sprayers

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage provide the most effective disease control. For ground application equipment, 4 - 16 litres of water/100m² is recommended.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use. Use a pump with the capacity to: (1) maintain a minimum of 240 KPa at nozzles, and (2) provide sufficient agitation in the tank to keep the mixture in suspension (this requires recirculation of 10% of the tank volume per minute). Use jet agitators or a liquid sparge tube for vigorous agitation.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh screens at the nozzles. Check nozzle manufacturer's recommendations. For information on spray equipment and calibration, consult sprayer manufacturer's and/or provincial or territorial recommendations. For specific local directions and spray schedules, consult the provincial or territorial recommendations.

Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the DISARM 480 SC Fungicide to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the DISARM 480 SC Fungicide has completely and uniformly dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

SENSITIVE AREAS

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions

CONTROLLING DROPLET SIZE

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Field Sprayer (Broadcast and Ground boom) application: **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Spray Buffer Zones:

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop /Site	Spray Buffer Zones (metres) Required for the Protection of:			
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field Sprayer (Broadcast and Ground boom)	Turf	1	1	10	5

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

SPRAY PREPARATION

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

NOTE: When treatment with DISARM 480 SC Fungicide has been completed, further field irrigation over the treated area should be avoided for 24 hours to prevent washing the chemical off the turf.

DISARM 480 SC Fungicide provides control of diseases in turf. DISARM 480 SC Fungicide should be used in conjunction with cultural practices that promote healthy, vigorous turf. These practices include nutrient management, thatch management, water management and judicious use of other fungicides and cultural practices.

Rate Ranges: Use the shorter specified application interval and/or the higher specified rate when prolonged favorable disease conditions exist.

DIRECTIONS FOR APPLICATION TO TURF

Disease Control	Use Rate (mL of product per 100 m ²)	App Interval (days)	Remarks
Anthracnose – Foliar Blight (<i>Colletotrichum cereale</i>)	5.8-11.5	14-28	Use preventatively. Begin applications when conditions are favourable for disease development. Under severe conditions, tank-mix with Daconil 2787 Flowable Fungicide at labelled rates.
Anthracnose -Basal Rot (<i>Colletotrichum cereale</i>)	11.5	14-21	Use preventatively. Begin applications when conditions are favourable for disease development. Under severe conditions, tank-mix with Daconil 2787 Flowable Fungicide at labelled rates.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	5.00 - 11.5	14 - 21	DISARM 480 SC Fungicide provides control of light to moderate dollar spot pressure. Under heavy dollar spot pressure use a Group 3 (DMI) fungicide labeled for control of Dollar Spot with DISARM 480 SC Fungicide applications.
Pink Snow Mould (<i>Microdochium nivale</i>)	11.5	n/a	Apply 1 application prior to permanent snow cover and before turf dormancy. Tank-mix with Daconil 2787 Flowable Fungicide at labelled rates to improve efficacy against grey snow mould caused by <i>Typhula ishikariensis</i> .
Grey Snow Mould (<i>Typhula incarnata</i>)	11.5	n/a	
Summer Patch (<i>Magnaporthe poae</i>)	5.8- 11.5	14-28	Start applications in the spring when the soil temperatures at 5 cm depth reach 15.5-18 °C or as prescribed by local turf specialists.

Take All Patch (<i>Gaeumannomyces graminis</i>)	11.5	28	Begin applications before disease is present and continue applications while conditions are favourable for disease development. Make up to 2 applications in the spring and/or fall. Maximum 3 applications per season.
Pythium Foliar Blight (<i>Pythium aphanidermatum</i>)	11.5	7-14	Use preventatively. Begin applications when conditions are favourable for disease infection, prior to disease symptom development. During periods of prolonged favourable conditions, treat on the 14-day application interval. Maximum 3 applications per season.
Brown Ring Patch (Waitea Patch) (<i>Waitea circinata</i> var. <i>circinata</i>)	5.8-11.5	14-21	Use preventatively. Begin applications when conditions are favourable for disease infection, prior to disease symptom development. Use the higher rate under high disease pressure.
Melting out and Leaf Spot (<i>Bipolaris sorokiniana</i> , <i>Drechslera poae</i>)	5.8-11.5	14-21	Apply when conditions are favourable for disease development. Applications should be made preventatively. The high rate should be used for curative applications, when disease pressure is high and/or when environmental conditions are favourable for disease development.
Brown Patch (<i>Rhizoctonia solani</i>)	5.8-11.5	14-28	Apply when conditions are favourable for disease development.

For control of pythium blight and anthracnose, do not apply more than 1 application of Disarm 480 SC Fungicide before alternating with a fungicide having a different mode of action registered for the same disease or pathogen. For control of diseases other than pythium blight and anthracnose, do not apply more than 2 sequential applications before alternating with a fungicide having a different mode of action registered for the same disease or pathogen.

Tank Mixing

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact [insert registrant name] at [insert contact information] for information before applying any tank mix that is not specifically recommended on this label.

Restrictions and Other Information

- Do not apply more than 40 mL/100m² of DISARM 480 SC Fungicide per year, or more than 11.5 mL/100m² per application.
- There is a maximum number of 4 applications per season.
- Not for home owner use. May only be applied to residential turf by a certified pest control operator.
- Under conditions of high disease pressure, use the higher rate, the shortest application interval or both.
- DISARM 480 SC Fungicide may be tank mixed at table rates to increase the spectrum of controlled diseases.

UPL, the UPL logo and DISARM are registered trademarks of a UPL Corporation Limited Group Company.
©2021 UPL Corporation Limited Group Company.